





# **Indian Journal of Postgraduate Dermatology**

Article in Press

#### Ouiz

# Syphilis Quiz

Abhinav Bansal<sup>1</sup>, Ananta Khurana<sup>1</sup>

Department of Dermatology, Venereology and Leprosy, Atal Bihari Vajpayee Institute of Medical Sciences and Dr. Ram Manohar Lohia Hospital, New Delhi,

#### \*Corresponding author:

Abhinav Bansal, Department of Dermatology, Venereology and Leprosy, Atal Bihari Vajpayee Institute of Medical Sciences and Dr Ram Manohar Lohia Hospital,

abhinavbansal866@gmail.com

New Delhi, India.

Received: 19 July 2024 Accepted: 12 September 2024 EPub Ahead of Print: 17 October 2024 Published:

#### DOI

10.25259/IJPGD\_157\_2024

#### **Quick Response Code:**



## **QUESTIONS**

- In 1927, Jauregg was awarded the Nobel Prize for Physiology and Medicine for which discovery?
  - a) Discovery of penicillin
  - b) Induction of malaria in syphilis patients
  - Discovery of Treponema pallidum
  - Discovery of Arsphenamine
- What is popularly known as 'magic bullet' or 'compound 606' in context of syphilis treatment?
  - Salvarsan
  - Penicillin
  - c) Bismuth salts
  - d) Mercurous chloride
- Which syphilis law states that a seemingly healthy baby born to a mother with syphilis cannot contract the infection from her mother through breastfeeding or from a wet nurse?
  - Colles law
  - b) Profeta law
  - Kassowitz law c)
  - Baumis law
- What is Dubois sign in congenital syphilis?
  - Cicatricial lines radiating from mouth
  - Bony prominence of forehead
  - Fracture through degenerated metaphysis
  - Short incurved little finger
- The earliest test to become positive for the detection of syphilis is
  - a) Venereal disease research laboratory (VDRL)
  - b) TPHA
  - c) Enzyme immunoassay
  - d) Treponema pallidum passive particle agglutination
- Which among the following is a cause of acute biological false positive VDRL
  - a) Leprosy
  - b) Malignancy

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms. ©2024 Published by Scientific Scholar on behalf of Indian Journal of Postgraduate Dermatology

- c) Pregnancy
- d) Old age
- 7. Patterns of alopecia seen in patients of secondary syphilis are all except?
  - a) Moth-eaten alopecia
  - b) Trichotilomania like
  - c) Alopecia areata like
  - d) Cicatricial alopecia
- Reappearance of gumma at the site of healed chancre is known as
  - a) Chancre redux
  - Pseudochancre redux
  - c) Mixed chancre
  - d) Kissing chancre
- The lesions over the palms as shown in Figure 1 are a feature of:
  - a) Primary syphilis
  - Secondary syphilis
  - c) Gumma
  - Condyloma lata d)



Figure 1: Multiple scaly copper penny rash over both palms.

- 10. According to the Centre for Disease Control and Prevention guidelines, what should be the acceptable missed dose interval in non-pregnant patients?
  - a) 10-14 days
  - b) 7-9 days
  - c) 10-11 days
  - d) 12-14 days
- 11. All of the following are the effect of human immunodeficiency virus on syphilis except?
  - a) Lues maligna
  - b) Multiple chancre
  - Delayed progression to tertiary stage neurosyphilis

- d) More relapse and recurrences
- 12. Who discovered Treponema pallidum?
  - Schaudinn and Hoffmann
  - b) Ricord
  - c) Landsteiner
  - d) Paracelsus
- 13. All are the advantages of dark ground microscopy except?
  - a) It provides immediate diagnosis
  - b) It can distinguish from other spiral organisms
  - Can diagnose syphilis even before serological tests become positive
  - It can be used for both oral and rectal lesions
- 14. Which of the following is not a non-treponemal test used to detect syphilis?
  - RPR a)
  - b) TRUST
  - c) USR
  - d) FTA-Abs test
- 15. Which of the following is not a part of stigmata of congenital syphilis?
  - Wimberger sign
  - Higoumenaki sign
  - Gothic palate
  - Frontal bossing
- 16. Partners who have had sexual intercourse within how many days with patients having syphilis should receive presumptive treatment for early syphilis?
  - 30 days a)
  - 45 days b)
  - 90 days
  - d) 60 days
- 17. The most common part of aorta affected in cardiovascular syphilis is?
  - a) Ascending aorta
  - b) Arch of aorta
  - Descending aorta c)
  - Abdominal aorta
- 18. Which of the morphological variant of secondary syphilis is typically itchy?
  - a) Maculopapular
  - b) Papulosquamous
  - c) Pustular
  - d) Follicular
- 19. Which of the following is not a complication of primary chancre?
  - Phagedenic chancre
  - Plaques fauchees en prairie
  - Thrombophlebitis of dorsal vein of penis c)
  - d) Syphilitic balanitis of Follman

- 20. Treatment failure to the syphilis treatment is suspected when
  - Patients with persistent or recurring signs/symptoms
  - Sustained 4-fold increase in non-treponemal titres at repeat testing
  - Failure of non-treponemal titres to decline 4-fold within 12 months
  - d) All of the above

#### **ANSWERS**

- 1. b
- 2. a
- 3. b
- 4. d
- 5. c
- 6. c
- 7. d
- 8. b
- 9. b
- 10. a
- 11. c
- 12. a
- 13. d
- 14. d
- 15. a
- 16. c
- 17. a
- 18. d

- 19. b
- 20. d

#### Ethical approval

Institutional Review Board approval is not required.

#### Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent.

### Financial support and sponsorship

Nil.

#### **Conflicts of interest**

Dr. Ananta Khurana is on the Editorial Board of the Journal.

# Use of artificial intelligence (AI)-assisted technology for manuscript preparation

The authors confirm that there was no use of artificial intelligence (AI)-assisted technology for assisting in the writing or editing of the manuscript and no images were manipulated using AI.

How to cite this article: Bansal A, Khurana A. Syphilis Quiz. Indian J Postgrad Dermatol. doi: 10.25259/IJPGD\_157\_2024