



Indian Journal of Postgraduate Dermatology



Quiz Syphilis Quiz

Abhinav Bansal¹, Ananta Khurana¹

¹Department of Dermatology, Venereology and Leprosy, Atal Bihari Vajpayee Institute of Medical Sciences and Dr. Ram Manohar Lohia Hospital, New Delhi, India.

*Corresponding author:

QUESTIONS

Abhinav Bansal, Department of Dermatology, Venereology and Leprosy, Atal Bihari Vajpayee Institute of Medical Sciences and Dr Ram Manohar Lohia Hospital, New Delhi, India.

abhinavbansal866@gmail.com

Received: 19 July 2024 Accepted: 12 September 2024 EPub Ahead of Print: 17 October 2024 Published: 07 February 2025

DOI

10.25259/IJPGD_157_2024

Quick Response Code:



- 1. In 1927, Jauregg was awarded the Nobel Prize for Physiology and Medicine for which discovery?
 - a) Discovery of penicillin
 - b) Induction of malaria in syphilis patients
 - c) Discovery of Treponema pallidum
 - d) Discovery of Arsphenamine
- 2. What is popularly known as 'magic bullet' or 'compound 606' in context of syphilis treatment?
 - a) Salvarsan
 - b) Penicillin
 - c) Bismuth salts
 - d) Mercurous chloride
- 3. Which syphilis law states that a seemingly healthy baby born to a mother with syphilis cannot contract the infection from her mother through breastfeeding or from a wet nurse?
 - a) Colles law
 - b) Profeta law
 - c) Kassowitz law
 - d) Baumis law
- 4. What is Dubois sign in congenital syphilis?
 - a) Cicatricial lines radiating from mouth
 - b) Bony prominence of forehead
 - c) Fracture through degenerated metaphysis
 - d) Short incurved little finger
- 5. The earliest test to become positive for the detection of syphilis is
 - a) Venereal disease research laboratory (VDRL)
 - b) TPHA
 - c) Enzyme immunoassay
 - d) Treponema pallidum passive particle agglutination
- 6. Which among the following is a cause of acute biological false positive VDRL
 - a) Leprosy
 - b) Malignancy

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms. ©2025 Published by Scientific Scholar on behalf of Indian Journal of Postgraduate Dermatology

- c) Pregnancy
- d) Old age
- 7. Patterns of alopecia seen in patients of secondary syphilis are all except?
 - a) Moth-eaten alopecia
 - b) Trichotilomania like
 - c) Alopecia areata like
 - d) Cicatricial alopecia
- 8. Reappearance of gumma at the site of healed chancre is known as
 - a) Chancre redux
 - b) Pseudochancre redux
 - c) Mixed chancre
 - d) Kissing chancre
- 9. The lesions over the palms as shown in Figure 1 are a feature of:
 - a) Primary syphilis
 - b) Secondary syphilis
 - c) Gumma
 - d) Condyloma lata



Figure 1: Multiple scaly copper penny rash over both palms.

- 10. According to the Centre for Disease Control and Prevention guidelines, what should be the acceptable missed dose interval in non-pregnant patients?
 - a) 10–14 days
 - b) 7-9 days
 - c) 10-11 days
 - d) 12-14 days
- 11. All of the following are the effect of human immunodeficiency virus on syphilis except?
 - a) Lues maligna
 - b) Multiple chancre
 - c) Delayed progression to tertiary stage and neurosyphilis

- d) More relapse and recurrences
- 12. Who discovered Treponema pallidum?
 - a) Schaudinn and Hoffmann
 - b) Ricord
 - c) Landsteiner
 - d) Paracelsus
- 13. All are the advantages of dark ground microscopy except?
 - a) It provides immediate diagnosis
 - b) It can distinguish from other spiral organisms
 - c) Can diagnose syphilis even before serological tests become positive
 - d) It can be used for both oral and rectal lesions
- 14. Which of the following is not a non-treponemal test used to detect syphilis?
 - a) RPR
 - b) TRUST
 - c) USR
 - d) FTA-Abs test
- 15. Which of the following is not a part of stigmata of congenital syphilis?
 - a) Wimberger sign
 - b) Higoumenaki sign
 - c) Gothic palate
 - d) Frontal bossing
- 16. Partners who have had sexual intercourse within how many days with patients having syphilis should receive presumptive treatment for early syphilis?
 - a) 30 days
 - b) 45 days
 - c) 90 days
 - d) 60 days
- 17. The most common part of aorta affected in cardiovascular syphilis is?
 - a) Ascending aorta
 - b) Arch of aorta
 - c) Descending aorta
 - d) Abdominal aorta
- 18. Which of the morphological variant of secondary syphilis is typically itchy?
 - a) Maculopapular
 - b) Papulosquamous
 - c) Pustular
 - d) Follicular
- 19. Which of the following is not a complication of primary chancre?
 - a) Phagedenic chancre
 - b) Plaques fauchees en prairie
 - c) Thrombophlebitis of dorsal vein of penis
 - d) Syphilitic balanitis of Follman

13. d

14. d

15. a

16. c

17. a

18. d

19. b

20. d

- 20. Treatment failure to the syphilis treatment is suspected when
 - a) Patients with persistent or recurring signs/symptoms
 - b) Sustained 4-fold increase in non-treponemal titres at repeat testing
 - c) Failure of non-treponemal titres to decline 4-fold within 12 months
 - d) All of the above

ANSWERS

1. b

2. a

- 3. b
- 4. d
- 5. c
- 6. c
- 7. d
- 8. b
- 9. b
- 10. a 11. c
- 11. c 12. a

Ethical approval: Institutional Review Board approval is not required.

Declaration of patient consent: The authors certify that they have obtained all appropriate patient consent.

Financial support and sponsorship: Nil.

Conflicts of interest: Dr. Ananta Khurana is on the Editorial Board of the Journal.

Use of artificial intelligence (AI)-assisted technology for manuscript preparation: The authors confirm that there was no use of artificial intelligence (AI)-assisted technology for assisting in the writing or editing of the manuscript and no images were manipulated using AI.

How to cite this article: Bansal A, Khurana A. Syphilis Quiz. Indian J Postgrad Dermatol. 2025;3:107-9. doi: 10.25259/IJPGD_157_2024